

Effect of salinity and inoculation on growth, nitrogen fixation and nutrient uptake of *Vigna radiata* (L.) Wilczek

FAUZIA Y. HAFEEZ, ZAHOOR ASLAM and KAUSER A. MALIK
Nuclear Institute for Agriculture and Biology, Faisalabad, Pakistan

Received 2 February 1987. Revised September 1987

Key words: inoculation, N₂-fixation, salinity, *Vigna radiata*

Abstract

This study reports the effect of salinity and inoculation on growth, ion uptake and nitrogen fixation by *Vigna radiata*. A soil EC_e level of 7.5 dS m⁻¹ was quite detrimental causing about 60% decline in dry matter and grain yield of mungbean plants whereas a soil EC_e level of 10.0 dS m⁻¹ was almost toxic. In contrast most of the studied strains of *Rhizobium* were salt tolerant. Nevertheless, nodulation, nitrogen fixation and total nitrogen concentration in the plant was drastically affected at high salt concentration. A noticeable decline in acetylene reduction activity occurred when salinity level increased to 7.5 dS m⁻¹.

Introduction

Vigna radiata is an important pulse crop in Pakistan. Being a legume it could produce food rich in protein without addition of nitrogen fertilizers. Therefore, this crop is especially suitable in developing countries like Pakistan where availability of protein is insufficient and fertilizers are expensive.

Nodulation in *Vigna radiata* under field condition is very poor (Idris *et al.*, in press; Ramaswamy and Nair, 1965). It may be due to absence of *Rhizobium* in such soils. Alternatively other environmental factors such as salinity (Abdel Ghaffar *et al.*, 1982; Lauter *et al.*, 1981), high temperature and drought (Marshall, 1964; Mass and Hoffman, 1977; Vincent *et al.*, 1962) may effect the nodulation and nitrogen fixation of leguminous plants. For example a few studies (Bernstein and Ogata, 1966; Wilson, 1970) show that the effect of salinity on nitrogen fixation ability of soybean and alfalfa could be very detrimental. In fact pulse crops are reported not to nodulate on salt-affected land even though native rhizobia are known to be present (Bhardwaj, 1974). The major point to investigate is whether symbiotic nitrogen fixation is more sensitive to salinity than host plant growth.

This paper reports the effect of salinity on the growth, nitrogen fixation, yield and nutrient uptake in mungbean. In addition, *in vitro* salt tolerance of five strains of *Rhizobium* spp. was also assessed.

Materials and methods

The seeds of mungbean (cv. 20–121) were obtained from Mutation Breeding Division, Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad. Three experiments were conducted in this study. The effect of different salt concentrations on the host (Mungbean) was studied in experiment A and B and on the endophyte (*Rhizobium*) was studied in experiment C.

Experiment A

This experiment was conducted in sterilized sand flushed with nutrient solution contained in Leonard jars. Seeds were inoculated with locally prepared carrier (gamma irradiated, filter mud amended with heavy loam and sucrose) based *Rhizobium* inoculum at a dose of 20 g 100 g⁻¹ seeds. Inoculum was mixture of five strains (M-11, M-17,

TAL 441, TAL 420, and TAL 169). For ensuring firm association of *Rhizobium* to the seeds, the seeds were successively coated with gum arabica, *Rhizobium* inoculum and rock phosphate. The number of viable cells at sowing time were $1.7 \times 10^{10} \text{ g}^{-1}$ of inoculum and $7 \times 10^6 \text{ seed}^{-1}$. Sand cultures were flushed daily with 1/4 strength nitrogen free Hoagland solution. At first leaf stage, thinning was done to leave uniform seedlings in each jar. At this stage NaCl, CaCl₂, Na₂SO₄ and MgCl₂ were added in ratio of 4:5:10:1 to the flushing nutrient solution to produce the salt concentrations of 1.4, 5.0, 7.5 and 10.0 dS m⁻¹. Each treatment was replicated six times. Plants were grown for 30 days. At harvesting shoot and root dry weight were recorded and roots were studied for nodulation and nitrogen fixation as estimated by acetylene reduction technique (Hardy *et al.*, 1968). Nitrogenase activity was measured by incubating excised nodulated root systems in 200-ml plastic bottles tightly closed with screwed caps fitted with suba seals. A 10:90 acetylene air atmosphere was created inside the bottle. After incubation for 1 hour at room temperature the gas samples (100 μ l) were analysed on a gas chromatograph (Carlo-Erba Model 180) fitted with a 1 m \times 2 mm steel column filled with Porapak R (80–100 mesh) and a H₂ flame ionization detector (FID). Nitrogen was used as a carrier gas at a flow rate of 30 ml min⁻¹. Two controls, one bottle with C₂H₂ but without nodules and the other with nodules but without addition of acetylene were also included during each assay to check indigenous production of ethylene.

Experiment B

This experiment was conducted in soil in 8 kg capacity plastic buckets. The soil used in the experiment was sandy clay loam, non-saline (EC_e = 1.4 dS m⁻¹; pH 7.8) and had available N and P of 0.039 and 2 ppm respectively. Available nitrogen in soil was determined according to Bremner (1965). Available phosphorus in the soil was assayed according to the Olsen method (Watanabe and Olsen, 1965). Artificial salinization of the soil was achieved with NaCl, CaCl₂, Na₂SO₄ and MgCl₂ mixed in ratio of 4:5:10:1 to produce salin-

ity levels of EC_e of 1.4, 5.0, 7.5 and 10 dS m⁻¹ (Qureshi *et al.*, 1977). Urea (20 kg ha⁻¹) and KH₂PO₄ (60 kg ha⁻¹) were applied at rates of 0.17 g and 1.7 g per pot respectively. Ten seeds were sown in each plastic pot. The pots were not drained and water was given as needed to keep the soil around field capacity. Seeds were inoculated as in experiment A. Each treatment was replicated twelve times. At first leaf stage the seedlings were thinned out to four uniform plants, in each pot.

Plants were sampled for nodulation and acetylene reduction assay at preflowering, 33 days after sowing (DAS) and flowering (45 DAS) stages and for yield at 65 DAS. Oven dried plant samples as well as air dried grains were ground and thoroughly homogenized. Such samples were used for nitrogen, phosphorus, potassium and sodium assay. For the estimation of total nitrogen concentration in plant, samples were digested in concentrated H₂SO₄ and measured by micro-Kjeldahl method (Bremner, 1965). For phosphorus, sodium and potassium assay, plant material was digested in a mixture of HNO₃ and HClO₄ (Richards *et al.*, 1954). Phosphorus concentration in plant was determined colorimetrically (Jackson, 1962). Potassium and sodium concentrations were measured by flame photometry. Each value was the mean of three replicates.

Experiment C

The salt tolerance of five *Rhizobium* strains (Cowpea group) was studied in yeast mannitol (Vincent, 1970) cultures containing 1.7, 5, 10, 25, 50, 100, 150 and 200 mol m⁻³ NaCl. The strains were chosen on the basis of their efficiency to nodulate mungbean. Several strains were studied and most effective were included in the experiment for further studies. Cells were grown for three days in the dark at 29 °C in shaken Erlenmeyer flask containing 50 ml medium. The initial cell density was 10⁶ viable cell ml⁻¹. Among the five *Rhizobium* strains, two strains (M-11, M-17) were of local origin while the rest (TAL 441, TAL 420, TAL 169) were obtained from NIFTAL Hawaii USA. Viable cells were counted by using standard serial dilution and plated by spread plate count method (Vincent, 1970).

Table 1. Effect of salinity and inoculation on dry matter yield of mungbean grown in soil. Plants were harvested at different stages of its growth (Experiment B)

| EC _c of soil dS m ⁻¹ | Dry weight (g plant ⁻¹) | | |
|--|-------------------------------------|-----------|----------|
| | Preflowering | Flowering | Maturity |
| 1.4 Uninoc. | 0.8 | 2.1b | 2.4ab |
| Inoc. | 1.1 | 2.7a | 3.1a |
| 5.0 Uninoc. | 0.6 | 1.0d | 1.9bc |
| Inoc. | 1.0 | 1.6c | 2.5ab |
| 7.5 Uninoc. | 0.4 | 0.6e | 1.0d |
| Inoc. | 0.6 | 1.1d | 1.3cd |
| 10.0 Uninoc. | 0.3 | 0.3e | |
| Inoc. | 0.3 | 0.4e | |
| NS | | | |

Means followed by same letter are not significantly different at 5% probability level.

Uninoc. Uninoculated.

Inoc. Inoculated.

NS: Non-significant.

Values are mean of three readings.

Results and discussion

Germination and plant survival

The germination was 100 percent in both soil and sand cultures at all salinity levels; only a delay of three days in germination was noted in the case of 7.5 and 10.0 dS m⁻¹. In fact in most plant species, moderate levels of salinity delay germination and not the germination percentage (Salim *et al.*, 1979). In experiment B the survival of plants was decreased with increasing salinity levels. The survival

Table 2. Effect of salinity and inoculation on dry matter yield and nodulation on mungbean grown in nutrient solution. Plants were harvested after 30 days of growth (Experiment A)

| EC _c dS m ⁻¹ | No. of nodule plant ⁻¹ | Frequency of nodulation (%) | Dry weight of nodules (mg plant ⁻¹) | μ mole C ₂ H ₄ g ⁻¹ dry nodule h ⁻¹ | Dr matter mg plant ⁻¹ (whole plant) |
|------------------------------------|-----------------------------------|-----------------------------|---|---|--|
| Uninoc. | | | | | |
| 1.4 | 0 | 0 | 0 | 0 | 753 ± 33 |
| Inoc. | | | | | |
| 1.4 | 7 | 85 | 18 ± 0.3 | 22 ± 2.2 | 901 ± 17 |
| 5.0 | 4 | 40 | 8 ± 0.2 | 18 ± 2.5 | 732 ± 13 |
| 7.5 | 2 | 10 | 4 ± 0.5 | 3 ± 1.6 | 417 ± 12 |
| 10.0 | 0 | 0 | 0 | 0 | 77 ± 6 |

Uninoc. Uninoculated.

Inoc. Inoculated.

Values are the mean of three readings ± standard deviation of the mean.

was 96% at 7.5 dS m⁻¹, at an EC_c level of 10 dS m⁻¹ the survival of plants were decreased from 80% to 58% at flowering stage and all plants had died by maturity. At high salt concentration death of seedlings after germination has also been reported in case of *Sesbania aculeata* (Salim *et al.*, 1979) and *Leucaena leucocephala* (Niaz *et al.*, 1985).

Plant growth and yield

Dry matter yield per plant (Table 1, 2) decreased significantly with increase in salinity levels regardless of the stages of plant growth. Data also showed that soil salinity of 7.5 dS m⁻¹ was quite detrimental to cause a 50–60% decline in dry matter yield of mungbean plant, whereas, the soil EC_c level of 10.0 dS m⁻¹ was almost toxic. Reduction in plant growth and dry matter accumulation was observed at moderate salinity levels in mungbean. This is in agreement with salt sensitivity reported in other leguminous plant species *Vicia faba*, *Phaseolus vulgaris* grown at the same salinity levels as reported in our experiments, (Abdel Ghaffar *et al.*, 1982) and *Glycine wightii* (Wilson, 1970).

An inverse correlation was found between salinity and grain weight. Higher the soil salinity the lower were the weight of grains (Table 3). However, the number of pods per plant and grains per pod were not significantly affected up to a salinity level of 7.5 dS m⁻¹. The grain yield of mungbean was about 60% lower at salinity levels of 5 dS m⁻¹ and 7.5 dS m⁻¹ as compared to 1.4 dS m⁻¹ while it was completely depressed at 10 dS m⁻¹. Similar results

Table 3. Effect of salinity and inoculation on some components of grain yield of mungbean grown in soil (Experiment B)

| EC _e of soil dS m ⁻¹ | Yield components | | No. of pod grain | | No. of grain 1000 grain | |
|--|------------------|-------|------------------|-------|-------------------------|-------|
| | Wt of | Wt of | pod | grain | Wt of | Wt of |
| 1.4 Uninoc. | 2.7 | 4.1 | 45.7a | 506a | 4.7a | 45.7a |
| 1.4 Inoc. | 4.0 | 4.7 | 43.4a | 816b | 4.7 | 43.4a |
| 5.0 Uninoc. | 2.3 | 3.7 | 25.2b | 214d | 3.7 | 25.2b |
| 5.0 Inoc. | 3.7 | 4.3 | 21.7b | 345c | 4.3 | 21.7b |
| 7.5 Uninoc. | 1.7 | 5.3 | 21.6b | 195d | 5.3 | 21.6b |
| 7.5 Inoc. | 2.3 | 5.3 | 27.0b | 330c | 5.3 | 27.0b |

Means followed by the same letter are not significantly different at 5% probability level.
 Uninoc. Uninoculated.
 Inoc. Inoculated.
 NS: Non-significant.
 Values are the mean of three readings.

have been reported on other leguminous plant species *Vicia*, *Phaseolus*, *Glycine* and *Medicago* (Abdel Ghaffar *et al.*, 1982; Bernstein and Ogata, 1966; Wilson, 1970).

Nodulation and nitrogen fixation

The response of mungbean to inoculation is shown in Table 2 and 4. In Experiment A, uninoculated plants bore no nodule and had a 16% lower dry matter yield than the inoculated plants at salinity level of EC_e 1.4 dS m⁻¹. In case of Experiment B, not a single nodule was observed in uninoculated plants, therefore, soil is either devoid of *Rhizobium* or indicates the ineffectiveness of local strains to infect mungbean roots. Similar results were obtained by Idri *et al.* (in press) in field grown mung-

bean plants. The frequency of nodulation on mungbean at flowering stage was 88%. In this experiment inoculation with mixture of five *Rhizobium* strains clearly benefited *V. radiata* as indicated with increased nodulation, nitrogen fixation, dry matter production and grain weight per plant. However, the effect of inoculation was not significant on the number of pods per plant and number of grains per pod as compared to controls.

The effect of salinity on nodulation and nitrogen fixation in mungbean is presented in Table 3 and 4. Nodulation and nitrogen fixation of inoculated plants were highly sensitive to salt. Frequency of nodulation, weight of nodules and nitrogen fixation was decreased by increasing salinization. Nodulation was reduced to about half at salinity level of EC_e 5.0 dS m⁻¹ as compared to 1.4 dS m⁻¹. The nodulation was completely depressed at EC_e 10 dS m⁻¹ regardless of plant growth stages. The effect of salinity was more severe on the number of nodules per plant than on the specific nitrogenase activity. A notable decline in acetylene reduction activity occurred when salinity level increased to 7.5 dS m⁻¹. These results corroborate the earlier findings on *Glycine*, *Macroptilium*, *Neonotonia*, *Medicago*, *Vicia* and *Phaseolus* species (Abdel Ghaffar *et al.*, 1982; Bernstein and Ogata, 1966; Lakshmi *et al.*, 1974; Wilson, 1970, 1985).

Nutrient uptake

The concentration of nitrogen in plant tissue as affected by soil salinity was in harmony with those of dry matter at maturity stage (Table 5). With the rise of salinization the total nitrogen concentration in pod grain and whole plant significantly decreased. The concentration of nitrogen significantly

| EC _e of soil dS m ⁻¹ | No. of nod. plant ⁻¹ | Frequency of nodulation (%) | Dry weight of nodule (mg) | Pre-flowering | | Flowering | |
|--|---------------------------------|-----------------------------|---------------------------|---------------|-----------|---------------|-----------|
| | | | | Pre-flowering | Flowering | Pre-flowering | Flowering |
| 1.4 | 14 | 20 | 66 | 88 | 12 ± 2.0 | 15 ± 5.4 | 38 ± 2.4 |
| 5.0 | 4 | 9 | 33 | 38 | 8 ± 2.1 | 11 ± 2.3 | 35 ± 1.3 |
| 7.5 | 2 | 1 | 33 | 8 | 4 ± 1.1 | 5 ± 2.0 | 23 ± 1.4 |
| 10.0 | 0 | 0 | 0 | 0 | - | - | 4 ± 1.5 |

Table 4. Effect of salinity and inoculation^a on nodulation and nitrogen fixation of mungbean grown in soil (Experiment B)

^a Uninoculated plants had no nodules. Therefore, data are given only for inoculated plants. Values are the mean of three readings.

Table 5. Effect of salinity and inoculation on total nitrogen concentration^a at maturity (Experiment B)

| EC _e of soil dSm ⁻¹ | Pod shell | Grains | Whole plant |
|--|-----------|--------|----------------|
| 1.4 Uninoc. | 11.1ab | 27.9b | 50.7bc |
| Inoc. | 13.6a | 32.9a | 59.4a |
| 5.0 Uninoc. | 8.1bc | 24.7cd | 45.6cd |
| Inoc. | 9.2abc | 26.4bc | 53.0ab |
| 7.5 Uninoc. | 4.7c | 23.0d | 44.1bcd |
| Inoc. | 5.0c | 26.3bc | 50.1d |

Means followed by the same letter are not significantly different at 5% probability level.

Uninoc. Uninoculated.

Inoc. Inoculated.

^a mg N g⁻¹ dry matter.

Values are the mean of three readings.

increased in grains and whole plant at all levels of salinity when inoculated with *Rhizobium*. The results are in agreement with those reported earlier on *Phaseolus vulgaris*. Inoculation of *P. vulgaris* markedly enhanced nodulation, N_2 -fixation plant dry matter N content and final yield (Abdel Ghaffar *et al.*, 1982). The P concentration was similar to N concentration in plant (data not presented) so is not discussed separately.

In general K^+ concentration was similar in different plant parts at preflowering and flowering stage of growth of mungbean grown at various salinity levels. In contrast higher the EC_e level in the soil, higher was the concentration of Na^+ in the shoot and plant as a whole at both preflowering and flowering stage. However, there was only a slight increase in concentration of Na^+ in pods and grain with increase in EC_e level of the soil (Table 6). The ions to the pods and grains have been mainly supplied through phloem. Therefore, the exclusion of Na^+ in the pods and grains may have been due to efficient exclusion of Na^+ in phloem vessels.

Experiment C

In this experiment five Bradyrhizobium strains 'Cowpea group' were checked for their relative salt tolerance. Out of these five strains, two strains M-11, M-17 were local isolates while TAL 441, TAL 420 and TAL 169 were obtained from NIF-TAL, Hawaii, USA. There was no significant difference in the growth of all the strains tested as even

Table 6. Effect of salinity on concentration^a of Na^+ and K^+ in mungbean grown in soil. Plants were harvested at different stages of growth (Experiment B)

| EC _e of soil dSm ⁻¹ | Flowering | | Maturity | | | | | |
|---|-----------|-------|-----------|-------|--------|-------|--------|-------|
| | | | Pod shell | | Grain | | Shoot | |
| | Na^+ | K^+ | Na^+ | K^+ | Na^+ | K^+ | Na^+ | K^+ |
| 1.4 | 23.9 | 85.7 | 4.8 | 77.4 | 3.6 | 66.4 | 28.5 | 83.8 |
| 5.0 | 23.0 | 86.3 | 5.8 | 58.6 | 4.6 | — | 43.0 | 86.4 |
| 7.5 | 50.0 | 85.0 | 7.6 | 75.4 | 5.9 | 67.0 | 61.0 | 88.0 |
| 10.0 | 75.6 | 76.4 | — | — | — | — | — | — |

^a meq 100 g⁻¹ dry matter.

Values are the mean of three readings.

200 mol m⁻³ NaCl did not effect the growth. Strain M-17 however showed a slight decrease (13%) in its cell numbers at 200 mol m⁻³ NaCl. Similar results showing relatively high salt tolerance of *Rhizobium* strains have also been reported by other workers while working with *Rhizobium meliloti* (Douka *et al.*, 1984; Kassem *et al.*, 1985; Singleton *et al.*, 1982).

This study suggests that salinity has an indirect effect on biological nitrogen fixation in mungbean. The rhizobia are generally more capable to cope with salinity than their host legumes. However, the effect of salinity was more pronounced on the number and weight of nodules per plant than on their specific nitrogenase activity. It is clear from Table 4 that at preflowering stage as a result of the increase in salinity, a reduction of 75% in nodule dry weight was estimated as compared to only 40% in case of nitrogenase activity. These observations indicate that in addition to indirect effects, salinity affects nodule formation. However, it seems that when a nodule is formed, then subsequently there is little influence of salinity on its functioning provided the plant can maintain a reasonable photosynthetic activity as in evident from the biomass yields at different salinity levels and the physiological age of the plant. More work is needed to explore the reasons why *Rhizobium* species fail to successfully infect mungbean roots at high salt concentrations in spite of their high *in vitro* tolerance to NaCl.

References

- Abdel-Ghaffar A S, El-Attar H A, El-Halfawi M H and Abdel-Salam A A 1982 Effect of inoculation, nitrogen fertilizer,

- salinity and water stress on symbiotic N₂-fixation by *Vicia faba* and *Phaseolus vulgaris*. In *Biological Nitrogen Fixation Technology for Tropical Agriculture*. Eds. P H Graham and S C Harris. pp 153-160. Centro International de Agricultura Tropical Cali, Colombia.
- Abdel-Rahman A M and Abdel Hadi A H 1984 Possibilities of reducing the adverse effects of salinity by IAA. *Biol. Plant (Prague)* 26, 81-87.
- Bernstein L and Ogata G 1966 Effects of salinity on nodulation, nitrogen fixation and growth of soybean and alfalfa. *Agron. J.* 58, 201-203.
- Bhardwaj K K 1974 Growth and symbiotic effectiveness of indigenous *Rhizobium* species in a saline-alkaline soil. *Proc. Indian Acad. Sci.* 40, 540-543.
- Bremner J M 1965 Total nitrogen and inorganic forms of nitrogen. In *Methods of Soil Analyses*. Ed. C A Black. pp 1149-1237. Vol. 2 Am. Soc. Agron. Medison, Wisconsin, USA.
- Douka C E, Xenoulis A C and Paradellis T 1984 Salinity tolerance of a *Rhizobium meliloti* strain isolated from salt affected soils. *Folia Microbiol.* 29, 316-324.
- Hardy R W F, Holsten R D, Jackson E K and Burns R C 1968 The acetylene ethylene assay for N₂-fixation: laboratory and field evaluation. *Plant Physiol.* 43, 1185-1207.
- Idris M, Ashraf M and Malik K A Response of mungbean (*Vigna radiata*) to *Rhizobium* inoculation for effective nodulation and nitrogen fixation under field conditions. *Pak. J. Soil Science (in press)*.
- Jackson M L 1962 *Soil chemical analysis*. Prentice Hall Inc, Englewood Cliffs, NJ p. 151.
- Kassem M, Capellano A and Gounot A M 1985 Effects du chlorure de sodium sur la croissance *in vitro*, l'inféctivité et l'efficacité de *Rhizobium meliloti*. *Mircen J. Appl. Microbiol. Biotech.* 1, 63-75.
- Lauter D J, Munns D N and Clarkin K L 1981 Salt response of chickpea as influenced by nitrogen supply. *Agron. J.* 73, 961-966.
- Lakshmi K M, Singh C S and Subba Rao N S 1974 Root hair infection and nodulation in lucerne (*Medicago sativa*) influenced by salinity and alkalinity. *Plant and Soil* 40, 261-268.
- Marshall K C 1964 Survival of root nodule bacteria in dry soils exposed to high temperature. *Aust. J. Agric. Res.* 15, 273-281.
- Mass E V and Hoffman G J 1977 Crop salt tolerance-current assessment. *J. Irrig. Drainage Div. ASCE* 103 (IR 2), 115-134.
- Niazi M L K, Haq M I and Malik K A 1985 Salt tolerance studies on ipil ipil (*Leucaena leucocephala* L) CV. K-8. *Pak. J. Bot.* 17, 43-47.
- Qureshi R H, Salim M, Aslam Z and Sandhu G R 1977 An improved gravel culture technique for salt tolerance studies on plant. *Pak. J. Agric. Sci.* 14, 11-18.
- Ramaswamy P P and Nair K S 1965 A study on the response to green gram to bacterial seed inoculation. *Madras Agri. J.* 52, 241-242.
- Richards L A 1968 *Diagnosis and improvement of saline and alkali soils*. Agriculture handbook No. 60 US Dept. Agric. Ind. Ed. Oxford and IBH Pub. Co. New Delhi, India.
- Salim M, Aslam Z, Sandhu G R and Qureshi R H 1979 Salt tolerance studies on *Sesbania aculeata*. *Pak. J. Agric. Sci.* XVI, 47-52.
- Singleton P W, El Swaify S A and Bohlool B B 1982 Effect of salinity on *Rhizobium* growth and survival. *Appl. Environ. Microbiol.* 44, 884-890.
- Vincent J M 1970 *A Manual for the Practical Study of the Root Nodule Bacteria*. Blackwell Scientific Publication, Oxford, England.
- Vincent J M, Thompson and Donovan K O 1962 Death of root nodule bacteria on drying. *Aust. J. Agric. Res.* 13, 258-270.
- Watanabe F S and Olsen S R 1965 Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil Sci. Soc. Am. Proc.* 29, 677-678.
- Wilson J R 1970 Response to salinity in Glycine. VI. Some effects of a range of short term salt stresses on the growth, nodulation and nitrogen fixation of *Glycine wightii*. *Aust. J. Agric. Res.* 21, 571-582.
- Wilson J R 1985 Comparative response to salinity of the growth and nodulation of *Macroptilium atropurpureum* cultivar Siratro and *Neonotonia wightii* cultivar Cooper seedlings. *Aust. J. Agric. Res.* 36, 589-600.