

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Canad. Math. Bull. Vol. XX (Y), ZZZZ pp. 1–8

Hamiltonian Properties of Generalized Halin Graphs

Shabnam Malik, Ahmad Mahmood Qureshi, and Tudor Zamfirescu

Abstract. A Halin graph is a graph $H = T \cup C$, where T is a tree with no vertex of degree two, and C is a cycle connecting the end-vertices of T in the cyclic order determined by a plane embedding of T . In this paper, we define classes of generalized Halin graphs, called k -Halin graphs, and investigate their Hamiltonian properties.

1 Introduction

A *Halin graph* is a graph H which is the union of a tree $T \neq K_2$ with no vertex of degree 2 and a cycle C connecting the end-vertices of T in the cyclic order determined by a plane embedding of T . Halin graphs are planar, 3-connected, and possess rather strong Hamiltonian properties. They are *1-Hamiltonian*, *i.e.*, they are Hamiltonian [2] and remain so after the removal of any single vertex, as Bondy showed (see [4]). Moreover, Barefoot proved that they are *Hamiltonian connected*, *i.e.*, they admit a Hamiltonian path between every pair of vertices [1]. Bondy and Lovász [3] and, independently, Skowrońska [6] proved that Halin graphs on n vertices are almost pancyclic; more precisely, they contain cycles of all lengths l ($3 \leq l \leq n$) except possibly for a single even length. Also, they showed that Halin graphs on n vertices whose vertices of degree 3 are all on the outer cycle C must be *pancyclic*, *i.e.*, they must contain cycles of all lengths from 3 to n , thus proving a conjecture of Malkevitch [5]. Can we generalize the notion of a Halin graph such that some of its Hamiltonian properties are preserved? In the present paper this is what we do. We generalize Halin graphs in the following way.

A 2-connected planar graph G without vertices of degree 2, possessing a cycle C such that

- (i) all vertices of C have degree 3 in G ,
- (ii) $G - C$ is connected and has at most k cycles

is called a k -Halin graph. The cycle C is called the *outer cycle* of G . The vertices and cycles in $G - C$ are called *inner vertices* and, respectively, *inner cycles* of G .

A 0-Halin graph is a usual Halin graph. Moreover, the class of k -Halin graphs is contained in the class of $(k + 1)$ -Halin graphs ($k \geq 0$). Thus we get a nested sequence of generalized Halin graphs. We shall see that, as expected, the Hamiltonicity of k -Halin graphs steadily decreases as k increases. Indeed, a 1-Halin graph is still

Received by the editors January 11, 2008.

AMS subject classification: Primary: 05C45; secondary: 05C38.

Keywords: k -Halin Graph, Hamiltonian, Hamiltonian connected, traceable.

©Canadian Mathematical Society ZZZZ.

Hamiltonian, but not necessarily Hamiltonian connected, a 2-Halin graph is not always Hamiltonian but still traceable, while a 3-Halin graph is not even necessarily traceable. The property of being 1-Hamiltonian, Hamiltonian connected or almost pancyclic is not preserved, even by 1-Halin graphs. However, Bondy and Lovász' result about the pancyclicity of Halin graphs with no inner vertex of degree 3 remains true even for 3-Halin graphs.

2 Hamiltonicity of 3-Halin Graphs

The graph obtained from a Halin graph by deleting a vertex x from its outer cycle is called a *reduced Halin graph* [3]. The three neighbouring vertices of x , whose degrees reduce by one, are called the *end-vertices* of the reduced Halin graph. Lemma 1 of [3] tells us the following.

Lemma 2.1 *In any reduced Halin graph each pair of end-vertices is joined by a Hamiltonian path.*

Lemma 2.1 will allow us to contract any reduced Halin subgraph of a graph G to a single vertex of degree 3, whenever we study Hamiltonian properties of G .

A path in a k -Halin graph will be called an *inner path*, if it has its end-vertices on distinct inner cycles and no other vertex on any inner or outer cycle.

We call a k -Halin graph ($k \geq 1$) *simple* if it is spanned by the union of all its inner paths and cycles and the outer cycle. Thus, a 1-Halin graph is simple if it has an inner cycle C_1 (besides the outer cycle C), and is spanned by $C \cup C_1$.

Theorem 2.2 *Every 1-Halin graph is Hamiltonian.*

Proof If the 1-Halin graph is also Halin, then it is Hamiltonian. Now let G be a 1-Halin graph with C_1 and C as its inner and outer cycles respectively (see Figure 1).




Figure 1

Let a be a vertex on C_1 and let $b \notin C_1$ be a neighbour of a . If $b \notin C$, the union of all paths from b to C , which do not contain a , is a tree T_b . This tree plus the edges on C between its leaves defines a reduced Halin graph H_b . We replace H_b by a single vertex $b' \in C$, adjacent with $a \in C_1$. If $b \in C$, we keep the edge ab . After doing

Hamiltonian Properties of Generalized Halin Graphs

3

this with all vertices of C_1 , G reduces to a simple 1-Halin graph consisting of the two cycles C and C_1 , and of edges between the two cycles, such that the outer cycle has only vertices of degree 3 (see Figure 2). A Hamiltonian cycle in this graph is shown in Figure 2. ■




Figure 2

Remark. A 1-Halin graph is not necessarily Hamiltonian connected. Indeed, Figure 3 shows a bipartite 1-Halin graph G with 4 black and 4 white vertices. A path between any pair of white vertices will have one more white vertex than black, so it cannot be Hamiltonian. ■




Figure 3: 1-Halin graph

A 2-Halin graph is not necessarily Hamiltonian. Indeed, Figure 4 shows a bipartite 2-Halin graph on 15 vertices. Such a graph has no Hamiltonian cycle.

Recall that a graph admitting a spanning path is called *traceable*, and the path is called *Hamiltonian*.

Theorem 2.3 *Every 2-Halin graph is traceable.*

Proof If the 2-Halin graph is also 1-Halin, then, by Theorem 2.2, it is Hamiltonian, hence traceable. Now let G be a 2-Halin graph with inner cycles C_1 and C_2 and outer cycle C , as shown in Figure 5.

Lemma 2.1 allows us to reduce G to a simple 2-Halin graph, that is, the union of C , C_1 , C_2 , and the unique path P between C_1 and C_2 in $G - C$ (possibly reduced




Figure 4: 2-Halin graph




Figure 5

to a vertex), plus edges between C and $C_1 \cup C_2 \cup P$ (see Figure 6). Let $a_1 \in C_1$ and $a_2 \in C_2$ be the endpoints of P . We claim that there is a Hamiltonian path in G between the neighbour b_1 or c_1 of a_1 on C_1 and the neighbour b_2 or c_2 of a_2 on C_2 . This is illustrated in Figure 6, where a path between b_1 and b_2 is realized. Accordingly, G is traceable. ■

Remark. A 3-Halin graph is not necessarily traceable. Indeed, Figure 7 shows a 3-Halin bipartite graph G with 22 vertices coloured in two colours, 12 black and 10 white.

3 Pancyclicity of 3-Halin Graphs

A graph on n vertices is called *almost pancyclic*, if it contains cycles of all lengths from 3 to n except possibly for one single length. Let us call *m-almost pancyclic* an almost pancyclic graph without cycles of length m .

As announced in the Introduction, we show here that all 3-Halin graphs without inner vertices of degree 3 are pancyclic, thus extending the corresponding result of Bondy and Lovász [3] on Halin graphs. We shall make use of the following central result of [3].

Hamiltonian Properties of Generalized Halin Graphs




Figure 6




Figure 7: 3-Halin graph

Lemma 3.1 *Every Halin graph is almost pancyclic. If the Halin graph H is m -almost pancyclic, then m is even and H must contain one of the three types of subgraphs depicted in Figure 8.*




Figure 8: ($m = 12$)

Theorem 3.2 Every 3-Halin graph without inner vertices of degree 3 is pancyclic.

Proof Let G be a 3-Halin graph without inner vertices of degree 3. There are at most 3 inner cycles in G . Choose an edge in each of them, such that no pair of edges has a common point. The total number of chosen edges can be two if the union of the 3 inner cycles is a cycle plus a chord. Delete these chosen edges. The resulting Halin graph H has at most 6 inner vertices of degree 3.

By Lemma 3.1, H is almost pancyclic. Assume cycles of length m are missing. Then, by Lemma 3.1, m is even and H must contain a reduced Halin graph of one of the types I, II, or III (Figure 8).

Suppose first that $m = 4$. Then H must contain a reduced Halin graph H' as described in Figure 9.




Figure 9

The point x of H' has degree 3. Hence it must belong in G to an edge e which has been deleted to obtain H . If the other endpoint of e is a vertex like x , i.e., a non-endpoint of a subgraph of H isomorphic to H' , then G has a cycle of length 4, and is therefore pancyclic. So, assume that the other endpoint of e is not a vertex like x . Since there are at most 3 edges like e , there are at most 3 vertices like x . But 4-almost pancyclic Halin graphs (see Figure 10) have more than 3 vertices like x if they are different from the graph H'' of Figure 10. In case $H = H''$, the vertex o must, on one hand, have degree at least 4 in G , but can, on the other hand, be no endpoint of any further edge of G . Thus, in any case we obtain a contradiction.




Figure 10

Hamiltonian Properties of Generalized Halin Graphs

Suppose now that $m = 6$. The smallest 6-almost pancyclic Halin graph is shown in Figure 11. This graph has 8 inner vertices of degree 3, so it cannot be H .




Figure 11

If $m = 8$, then, by Lemma 3.1, H must contain one of the reduced Halin subgraphs of Figure 12. Thus H has at least 6 inner vertices of degree 3, but they cannot be endpoints of only 3 edges in G , excepting the cases shown in Figure 13. In these cases, however, G has cycles of length 8, and is therefore pancyclic.




Figure 12



Figure 13

If $m \geq 10$, then the reduced Halin graph which must, by Lemma 3.1, appear in H has at least 8 inner vertices of degree 3, which is impossible. ■

The 37-Halin graph of Figure 14 has no cycle of length 4 and shows that not every k -Halin graph with no inner vertex of degree 3 must be pancyclic. So we are led to the following question.




Figure 14

Which is the maximal number k for which every k -Halin graph with no inner vertex of degree 3 is pancyclic?

References

- [1] C. A. Barefoot, *Hamiltonian connectivity of the Halin graphs*. Congr. Numer. **58**(1987) 93–102.
- [2] J. A. Bondy, *Pancyclic graphs: recent results*. In: Infinite and Finite Sets. Colloq. Math. Soc. János Bolyai 10, North-Holland, Amsterdam, 1975, pp. 181–187.
- [3] J. A. Bondy and L. Lovász, *Lengths of cycles in Halin graphs*. J. Graph Theory **9**(1985) 397–410.
- [4] L. Lovász and M. D. Plummer, *On a family of planar bicritical graphs*. Proc. London Math. Soc. **30**(1975) 160–176.
- [5] J. Malkevitch, *Cycle lengths in polytopal graphs*. In: Theory and Applications of Graphs, Lectures Notes in Math. 642. Springer Berlin, 1978 pp. 364–370.
- [6] M. Skowrońska, *The pancyclicity of Halin graphs and their exterior contractions*, In: Cycles in Graphs, North-Holland Math. Stud. 115. North-Holland, Amsterdam, 1985, pp. 179–194.

Abdus Salam School of Mathematical Sciences, GC University, 68-B, New Muslim Town, Lahore, Pakistan
 e-mail: shabnam.malik@gmail.com
 sirahmad@gmail.com

Faculty of Mathematics, University of Dortmund, 44221 Dortmund, Germany,

and

Institute of Mathematics, Romanian Academy, Bucharest, Romania
 e-mail: tzamfirescu@yahoo.com