


1-(5-Nitro-2-oxoindolin-3-ylidene)-4-*o*-tolylthio- semicarbazide methanol monosolvate

**Humayun Pervez, Muhammad Yaqub, Nazia Manzoor, M. Nawaz Tahir
and M. Saeed Iqbal**

Acta Cryst. (2009). **E65**, o2858

This open-access article is distributed under the terms of the Creative Commons Attribution Licence
<http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and
reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals Online is available from journals.iucr.org

$b = 7.6986(3)$ Å
 $c = 18.5937(6)$ Å
 $\beta = 119.847(2)^\circ$
 $V = 1769.07(11)$ Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.22$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.16 \times 0.12$ mm

1-(5-Nitro-2-oxoindolin-3-ylidene)-4-o-tolylthiosemicarbazide methanol monosolvate

Humayun Pervez,^a Muhammad Yaqub,^a Nazia Manzoor,^a M. Nawaz Tahir^{b*} and M. Saeed Iqbal^c

^aDepartment of Chemistry, Bahauddin Zakariya University, Multan 60800, Pakistan,

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan, and

^cDepartment of Chemistry, Government College University, Lahore, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 21 October 2009; accepted 22 October 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.002$ Å;
 R factor = 0.038; wR factor = 0.098; data-to-parameter ratio = 16.1.

In the title compound, $C_{16}H_{13}N_5O_3S\cdot CH_4O$, the dihedral angle between the isatin unit and the 2-methylphenyl group is $41.81(2)^\circ$ and intramolecular N—H···O and N—H···N hydrogen bonds occur, generating $S(6)$ and $S(5)$ rings, respectively. In the crystal, polymeric chains arise as a result of N—H···O, O—H···S and C—H···O interactions.

Related literature

For related structures, see: Revenko *et al.* (1994); Pervez *et al.* (2009). For graph-set theory, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$C_{16}H_{13}N_5O_3S\cdot CH_4O$
 $M_r = 387.42$

Monoclinic, $P2_1/c$
 $a = 14.2485(5)$ Å

Data collection

Bruker Kappa APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{min} = 0.963$, $T_{max} = 0.974$

18543 measured reflections
 4005 independent reflections
 2975 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.098$
 $S = 1.03$
 4005 reflections

249 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D\cdots H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3···O1	0.86	2.04	2.7074 (17)	134
N4—H4A···N2	0.86	2.20	2.6254 (18)	110
N1—H1···O4 ⁱ	0.86	2.02	2.8394 (19)	160
O4—H4B···S1 ⁱⁱ	0.82	2.55	3.3485 (14)	164
C16—H16C···O3 ⁱⁱⁱ	0.96	2.45	3.342 (3)	154

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 1, -y, -z + 1$; (iii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

We acknowledge partial funding of this research work and the award of an Indigenous Ph.D. scholarship to NM by the Higher Education Commission, Islamabad, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5167).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Pervez, H., Yaqub, M., Manzoor, N., Tahir, M. N. & Iqbal, M. S. (2009). *Acta Cryst. E65*, o2698–o2699.
- Revenko, M. D., Kravtsov, V. K. & Simonov, Yu. A. (1994). *Crystallogr. Rep.* **39**, 42–46.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o2858 [doi:10.1107/S1600536809043633]

1-(5-Nitro-2-oxoindolin-3-ylidene)-4-*o*-tolylthiosemicarbazide methanol monosolvate

H. Pervez, M. Yaqub, N. Manzoor, M. N. Tahir and M. S. Iqbal

Comment

Recently we have reported the preparation and crystal structure of (*Z*)-4-Hexyl-1-(5-nitro-2-oxo-2,3-dihydro-1H-indol-3-ylidene) thiosemicarbazide (Pervez *et al.*, 2009). The title compound (I, Fig. 1) has been prepared and being reported in continuation of synthesizing various isatin derivatives due to their importance.

The crystal structure of (II) Isatin β -4-(*p*-tolyl)thiosemicarbazone (Revenko *et al.*, 1994) has been published. The title compound (I) differs from (II) due to attachment of NO₂ group with isatin and positional change of CH₃ group on the benzene ring.

In the crystal structure of (I), the group A (C1—C8/N1/N2/N3/O1) of isatin moiety and 2-methylphenyl group B (C10—C17) are planar with a maximum r. m. s. deviations of 0.0187 and 0.0065 Å respectively, from their mean square plane. The dihedral angle between A/B is 41.81 (2) $^{\circ}$. The nitro group C (N2/O2/O3) is oriented at a dihedral angle of 5.7 (2) $^{\circ}$ with group A. In (I), there exist two intermolecular H-bondings resulting in S(5) and S(6) ring motifs (Bernstein *et al.*, 1995). Methanol monosolvate interlinks the molecules through H-bondings (Table 1., Fig. 2). The molecules are stabilized in the form of infinite one dimensional polymeric chains.

Experimental

4-*o*-Tolylthiosemicarbazide (0.45 g, 2.5 mmol) dissolved in ethanol (10 ml) was added to a hot solution of 5-nitroisatin (0.46 g, 2.5 mmol) in 50% aqueous ethanol (30 ml) containing a few drops of glacial acetic acid. The reaction mixture was then refluxed for 2 h. The yellow crystalline solid formed during heating under reflux was collected by suction filtration. Thorough washing with hot aqueous ethanol furnished the title compound (I) in pure form (0.71 g, 80%), m.p. 499 K. The yellow needles of (I) were grown in ethanol:n-hexane (1:4) system at room temperature by diffusion method.

Refinement

The H-atoms were positioned geometrically (O—H = 0.82 Å, N—H = 0.86 Å, C—H = 0.93–0.96 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures




Fig. 1. View of (I) with displacement ellipsoids drawn at the 50% probability level. H-atoms are shown by circles of arbitrary radius and the dotted lines represent the intramolecular H-bonds.

supplementary materials




Fig. 2. The partial packing of (I), which shows that molecules form infinite one dimensional polymeric chains.

1-(5-Nitro-2-oxoindolin-3-ylidene)-4-o-tolylthiosemicarbazide methanol monosolvate

Crystal data

C ₁₆ H ₁₃ N ₅ O ₃ S·CH ₄ O	$F_{000} = 808$
$M_r = 387.42$	$D_x = 1.455 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 2975 reflections
$a = 14.2485 (5) \text{ \AA}$	$\theta = 2.5\text{--}27.5^\circ$
$b = 7.6986 (3) \text{ \AA}$	$\mu = 0.22 \text{ mm}^{-1}$
$c = 18.5937 (6) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 119.847 (2)^\circ$	Cut needle, yellow
$V = 1769.07 (11) \text{ \AA}^3$	$0.30 \times 0.16 \times 0.12 \text{ mm}$
$Z = 4$	

Data collection

Bruker Kappa APEXII CCD diffractometer	4005 independent reflections
Radiation source: fine-focus sealed tube	2975 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.031$
Detector resolution: 7.60 pixels mm^{-1}	$\theta_{\text{max}} = 27.5^\circ$
$T = 296 \text{ K}$	$\theta_{\text{min}} = 2.5^\circ$
ω scans	$h = -18 \rightarrow 18$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$k = -10 \rightarrow 10$
$T_{\text{min}} = 0.963$, $T_{\text{max}} = 0.974$	$l = -23 \rightarrow 23$
18543 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained
$wR(F^2) = 0.098$	$w = 1/[\sigma^2(F_o^2) + (0.0413P)^2 + 0.4905P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
4005 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$

249 parameters

$\Delta\rho_{\min} = -0.22 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct
methods

Extinction coefficient: ?

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.36128 (4)	-0.18526 (6)	0.42815 (3)	0.0480 (2)
O1	0.43184 (10)	0.09444 (15)	0.66176 (7)	0.0471 (4)
O2	0.09516 (11)	0.88792 (17)	0.44105 (8)	0.0535 (5)
O3	0.14063 (12)	1.04229 (17)	0.54995 (9)	0.0631 (5)
N1	0.39348 (11)	0.35294 (18)	0.70502 (8)	0.0388 (4)
N2	0.28711 (10)	0.24879 (17)	0.49325 (8)	0.0342 (4)
N3	0.33161 (11)	0.09298 (17)	0.49406 (8)	0.0390 (4)
N4	0.20990 (10)	0.06337 (17)	0.35699 (8)	0.0364 (4)
N5	0.14422 (11)	0.90750 (18)	0.51621 (9)	0.0408 (5)
C1	0.28486 (11)	0.4853 (2)	0.58087 (9)	0.0309 (4)
C2	0.22200 (12)	0.6183 (2)	0.52973 (10)	0.0329 (5)
C3	0.21086 (12)	0.7648 (2)	0.56800 (10)	0.0342 (5)
C4	0.25938 (13)	0.7812 (2)	0.65330 (10)	0.0380 (5)
C5	0.32173 (13)	0.6488 (2)	0.70422 (10)	0.0385 (5)
C6	0.33402 (12)	0.5016 (2)	0.66721 (9)	0.0330 (4)
C7	0.38889 (12)	0.2370 (2)	0.64829 (10)	0.0362 (5)
C8	0.31690 (12)	0.3173 (2)	0.56487 (9)	0.0314 (4)
C9	0.29577 (12)	-0.0024 (2)	0.42330 (9)	0.0346 (5)
C10	0.14634 (12)	-0.0135 (2)	0.27704 (9)	0.0330 (5)
C11	0.11584 (12)	0.0923 (2)	0.20843 (10)	0.0367 (5)
C12	0.04962 (14)	0.0193 (3)	0.13070 (10)	0.0470 (6)
C13	0.01474 (14)	-0.1498 (3)	0.12170 (11)	0.0501 (6)
C14	0.04524 (13)	-0.2512 (2)	0.19058 (12)	0.0466 (6)
C15	0.11076 (14)	-0.1830 (2)	0.26893 (11)	0.0413 (5)
C16	0.15262 (16)	0.2771 (3)	0.21779 (12)	0.0543 (6)
O4	0.46564 (14)	0.2556 (2)	0.37038 (8)	0.0764 (6)
C17	0.43827 (17)	0.4240 (3)	0.37829 (13)	0.0599 (7)
H1	0.42879	0.33638	0.75775	0.0465*
H2	0.18875	0.60978	0.47231	0.0395*
H3	0.38341	0.05337	0.53999	0.0468*
H4	0.24971	0.88234	0.67621	0.0456*

supplementary materials

H4A	0.19014	0.16531	0.36307	0.0437*
H5	0.35450	0.65800	0.76159	0.0462*
H12	0.02844	0.08685	0.08361	0.0563*
H13	-0.02945	-0.19550	0.06905	0.0601*
H14	0.02187	-0.36585	0.18459	0.0559*
H15	0.13064	-0.25081	0.31575	0.0496*
H16A	0.23026	0.28089	0.24674	0.0814*
H16B	0.12630	0.34033	0.24866	0.0814*
H16C	0.12490	0.32864	0.16396	0.0814*
H4B	0.50868	0.21697	0.41631	0.0916*
H17A	0.38390	0.42215	0.39442	0.0898*
H17B	0.41066	0.48319	0.32618	0.0898*
H17C	0.50125	0.48363	0.41975	0.0898*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0570 (3)	0.0429 (3)	0.0355 (3)	0.0174 (2)	0.0165 (2)	-0.0014 (2)
O1	0.0506 (7)	0.0385 (7)	0.0345 (7)	0.0088 (5)	0.0077 (5)	0.0026 (5)
O2	0.0627 (8)	0.0480 (8)	0.0382 (8)	0.0117 (6)	0.0164 (6)	0.0063 (6)
O3	0.0910 (10)	0.0387 (8)	0.0612 (9)	0.0172 (7)	0.0390 (8)	-0.0016 (7)
N1	0.0443 (7)	0.0392 (8)	0.0220 (7)	-0.0009 (6)	0.0083 (6)	-0.0004 (6)
N2	0.0372 (7)	0.0302 (7)	0.0291 (7)	0.0014 (5)	0.0119 (6)	-0.0027 (6)
N3	0.0428 (7)	0.0361 (8)	0.0265 (7)	0.0084 (6)	0.0084 (6)	-0.0021 (6)
N4	0.0441 (7)	0.0288 (7)	0.0285 (7)	0.0046 (6)	0.0121 (6)	-0.0030 (6)
N5	0.0475 (8)	0.0347 (8)	0.0436 (9)	0.0026 (6)	0.0253 (7)	0.0013 (7)
C1	0.0333 (7)	0.0308 (8)	0.0254 (8)	-0.0043 (6)	0.0122 (6)	-0.0032 (6)
C2	0.0359 (8)	0.0346 (8)	0.0251 (8)	-0.0021 (6)	0.0129 (6)	-0.0012 (7)
C3	0.0361 (8)	0.0322 (8)	0.0341 (9)	-0.0005 (6)	0.0174 (7)	0.0012 (7)
C4	0.0428 (8)	0.0359 (9)	0.0364 (9)	-0.0039 (7)	0.0205 (7)	-0.0094 (7)
C5	0.0430 (9)	0.0437 (10)	0.0258 (8)	-0.0063 (7)	0.0149 (7)	-0.0077 (7)
C6	0.0346 (7)	0.0333 (8)	0.0269 (8)	-0.0052 (6)	0.0122 (6)	-0.0016 (7)
C7	0.0355 (8)	0.0348 (9)	0.0285 (8)	-0.0031 (7)	0.0085 (6)	-0.0006 (7)
C8	0.0325 (7)	0.0305 (8)	0.0254 (8)	-0.0024 (6)	0.0100 (6)	-0.0006 (7)
C9	0.0397 (8)	0.0339 (9)	0.0288 (8)	0.0004 (7)	0.0159 (7)	-0.0005 (7)
C10	0.0339 (7)	0.0347 (9)	0.0278 (8)	0.0030 (6)	0.0134 (6)	-0.0027 (7)
C11	0.0347 (8)	0.0391 (9)	0.0333 (9)	0.0045 (7)	0.0146 (7)	0.0024 (7)
C12	0.0433 (9)	0.0588 (12)	0.0279 (9)	0.0085 (8)	0.0095 (7)	0.0049 (8)
C13	0.0396 (9)	0.0605 (12)	0.0356 (10)	0.0019 (8)	0.0077 (8)	-0.0153 (9)
C14	0.0428 (9)	0.0401 (10)	0.0518 (11)	-0.0056 (8)	0.0196 (8)	-0.0132 (9)
C15	0.0472 (9)	0.0368 (9)	0.0390 (10)	-0.0003 (7)	0.0207 (8)	0.0001 (8)
C16	0.0581 (11)	0.0461 (11)	0.0468 (11)	-0.0016 (9)	0.0171 (9)	0.0114 (9)
O4	0.1094 (13)	0.0594 (9)	0.0329 (8)	0.0304 (9)	0.0147 (8)	-0.0028 (7)
C17	0.0589 (11)	0.0478 (12)	0.0609 (14)	0.0006 (9)	0.0206 (10)	-0.0080 (10)

Geometric parameters (\AA , $^\circ$)

S1—C9	1.6666 (17)	C5—C6	1.381 (2)
O1—C7	1.220 (2)	C7—C8	1.502 (2)

O2—N5	1.2215 (19)	C10—C11	1.389 (2)
O3—N5	1.227 (2)	C10—C15	1.381 (2)
O4—C17	1.383 (3)	C11—C16	1.496 (3)
O4—H4B	0.8200	C11—C12	1.392 (2)
N1—C7	1.358 (2)	C12—C13	1.374 (3)
N1—C6	1.389 (2)	C13—C14	1.373 (3)
N2—N3	1.353 (2)	C14—C15	1.384 (3)
N2—C8	1.292 (2)	C2—H2	0.9300
N3—C9	1.365 (2)	C4—H4	0.9300
N4—C10	1.428 (2)	C5—H5	0.9300
N4—C9	1.331 (2)	C12—H12	0.9300
N5—C3	1.458 (2)	C13—H13	0.9300
N1—H1	0.8600	C14—H14	0.9300
N3—H3	0.8600	C15—H15	0.9300
N4—H4A	0.8600	C16—H16C	0.9600
C1—C2	1.381 (2)	C16—H16A	0.9600
C1—C8	1.451 (2)	C16—H16B	0.9600
C1—C6	1.402 (2)	C17—H17A	0.9600
C2—C3	1.384 (2)	C17—H17B	0.9600
C3—C4	1.385 (2)	C17—H17C	0.9600
C4—C5	1.374 (2)		
C17—O4—H4B	109.00	N4—C10—C15	120.89 (14)
C6—N1—C7	111.53 (13)	N4—C10—C11	117.36 (14)
N3—N2—C8	116.02 (13)	C10—C11—C12	117.23 (16)
N2—N3—C9	121.18 (13)	C10—C11—C16	121.31 (15)
C9—N4—C10	128.37 (14)	C12—C11—C16	121.46 (16)
O2—N5—C3	118.45 (14)	C11—C12—C13	121.76 (17)
O3—N5—C3	118.61 (14)	C12—C13—C14	119.86 (17)
O2—N5—O3	122.94 (15)	C13—C14—C15	120.05 (16)
C6—N1—H1	124.00	C10—C15—C14	119.48 (15)
C7—N1—H1	124.00	C1—C2—H2	122.00
N2—N3—H3	119.00	C3—C2—H2	122.00
C9—N3—H3	119.00	C5—C4—H4	120.00
C10—N4—H4A	116.00	C3—C4—H4	120.00
C9—N4—H4A	116.00	C4—C5—H5	121.00
C2—C1—C6	120.36 (15)	C6—C5—H5	121.00
C2—C1—C8	133.02 (14)	C13—C12—H12	119.00
C6—C1—C8	106.61 (13)	C11—C12—H12	119.00
C1—C2—C3	116.82 (15)	C12—C13—H13	120.00
N5—C3—C4	118.52 (14)	C14—C13—H13	120.00
N5—C3—C2	118.56 (14)	C15—C14—H14	120.00
C2—C3—C4	122.92 (15)	C13—C14—H14	120.00
C3—C4—C5	120.31 (15)	C10—C15—H15	120.00
C4—C5—C6	117.66 (15)	C14—C15—H15	120.00
N1—C6—C1	109.71 (14)	C11—C16—H16B	109.00
N1—C6—C5	128.35 (14)	C11—C16—H16C	109.00
C1—C6—C5	121.94 (15)	C11—C16—H16A	109.00
N1—C7—C8	106.00 (13)	H16A—C16—H16C	109.00
O1—C7—C8	126.66 (15)	H16B—C16—H16C	109.00

supplementary materials


O1—C7—N1	127.32 (15)	H16A—C16—H16B	109.00
N2—C8—C1	126.86 (14)	O4—C17—H17A	109.00
C1—C8—C7	106.14 (13)	O4—C17—H17B	109.00
N2—C8—C7	126.98 (15)	O4—C17—H17C	109.00
N3—C9—N4	114.76 (14)	H17A—C17—H17B	109.00
S1—C9—N4	127.11 (12)	H17A—C17—H17C	109.00
S1—C9—N3	118.14 (12)	H17B—C17—H17C	109.00
C11—C10—C15	121.60 (15)		
C7—N1—C6—C1	1.2 (2)	C6—C1—C8—N2	178.05 (18)
C7—N1—C6—C5	-178.24 (19)	C6—C1—C8—C7	-0.3 (2)
C6—N1—C7—O1	-179.54 (19)	C1—C2—C3—N5	-179.37 (17)
C6—N1—C7—C8	-1.4 (2)	C1—C2—C3—C4	0.0 (3)
C8—N2—N3—C9	171.16 (17)	N5—C3—C4—C5	179.22 (18)
N3—N2—C8—C1	177.28 (17)	C2—C3—C4—C5	-0.2 (3)
N3—N2—C8—C7	-4.8 (3)	C3—C4—C5—C6	0.3 (3)
N2—N3—C9—S1	175.23 (13)	C4—C5—C6—N1	179.14 (19)
N2—N3—C9—N4	-5.1 (2)	C4—C5—C6—C1	-0.3 (3)
C10—N4—C9—S1	7.3 (3)	O1—C7—C8—N2	0.9 (3)
C10—N4—C9—N3	-172.36 (17)	O1—C7—C8—C1	179.19 (19)
C9—N4—C10—C11	-137.33 (19)	N1—C7—C8—N2	-177.33 (18)
C9—N4—C10—C15	47.0 (3)	N1—C7—C8—C1	1.0 (2)
O2—N5—C3—C2	4.6 (3)	N4—C10—C11—C12	-177.02 (18)
O2—N5—C3—C4	-174.76 (18)	N4—C10—C11—C16	2.9 (3)
O3—N5—C3—C2	-174.74 (18)	C15—C10—C11—C12	-1.4 (3)
O3—N5—C3—C4	5.9 (3)	C15—C10—C11—C16	178.6 (2)
C6—C1—C2—C3	0.0 (3)	N4—C10—C15—C14	177.10 (18)
C8—C1—C2—C3	-178.49 (19)	C11—C10—C15—C14	1.6 (3)
C2—C1—C6—N1	-179.38 (17)	C10—C11—C12—C13	0.6 (3)
C2—C1—C6—C5	0.1 (3)	C16—C11—C12—C13	-179.4 (2)
C8—C1—C6—N1	-0.5 (2)	C11—C12—C13—C14	0.0 (3)
C8—C1—C6—C5	178.98 (18)	C12—C13—C14—C15	0.2 (3)
C2—C1—C8—N2	-3.3 (3)	C13—C14—C15—C10	-1.0 (3)
C2—C1—C8—C7	178.38 (19)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N3—H3 \cdots O1	0.86	2.04	2.7074 (17)	134
N4—H4A \cdots N2	0.86	2.20	2.6254 (18)	110
N1—H1 \cdots O4 ⁱ	0.86	2.02	2.8394 (19)	160
O4—H4B \cdots S1 ⁱⁱ	0.82	2.55	3.3485 (14)	164
C16—H16C \cdots O3 ⁱⁱⁱ	0.96	2.45	3.342 (3)	154

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x+1, -y, -z+1$; (iii) $x, -y+3/2, z-1/2$.

Fig. 1



supplementary materials

Fig. 2

